

# Unit Interval Orders and Hessenberg Varieties

Timothy Y. Chow

Center for Communications Research

July 9, 2017

## Stanley's chromatic symmetric function $\chi_G$

Let  $G$  be a graph with vertex set  $V = \{1, 2, \dots, n\}$ .

A **coloring**  $\kappa$  of  $G$  is a map  $\kappa : V \rightarrow \mathbb{N}$  such that adjacent vertices are mapped to different integers (“colors”).

# Stanley's chromatic symmetric function $X_G$

Let  $G$  be a graph with vertex set  $V = \{1, 2, \dots, n\}$ .

A **coloring**  $\kappa$  of  $G$  is a map  $\kappa : V \rightarrow \mathbb{N}$  such that adjacent vertices are mapped to different integers (“colors”).

**Definition.** The **chromatic symmetric function**  $X_G$  is

$$X_G := \sum_{\kappa} x_{\kappa(1)} x_{\kappa(2)} \cdots x_{\kappa(n)},$$

where the sum is over all colorings  $\kappa$  of  $G$ .

## Stanley's chromatic symmetric function $X_G$

Let  $G$  be a graph with vertex set  $V = \{1, 2, \dots, n\}$ .


A **coloring**  $\kappa$  of  $G$  is a map  $\kappa : V \rightarrow \mathbb{N}$  such that adjacent vertices are mapped to different integers (“colors”).

**Definition.** The **chromatic symmetric function**  $X_G$  is

$$X_G := \sum_{\kappa} x_{\kappa(1)} x_{\kappa(2)} \cdots x_{\kappa(n)},$$

where the sum is over all colorings  $\kappa$  of  $G$ .

**Example.**



$$\begin{aligned} X_G &= 6(x_1 x_2 x_3 + x_1 x_2 x_4 + \cdots) + (x_1^2 x_2 + x_2^2 x_1 + x_1^2 x_3 + x_3^2 x_1 + \cdots) \\ &= 6e_3 + e_1 e_2 - 3e_3 \\ &= 3e_3 + e_1 e_2. \end{aligned}$$

# Indifference graphs

Note that in the above example,  $X_G$  is **e-positive**; i.e., it is a polynomial in the  $e_i$  with **nonnegative** coefficients.

This is not true for all graphs; however. . .

# Indifference graphs

Note that in the above example,  $X_G$  is **e-positive**; i.e., it is a polynomial in the  $e_i$  with **nonnegative** coefficients.

This is not true for all graphs; however. . .

**Definition.** An **indifference graph** is a graph whose vertex set is a set of (distinct) closed unit intervals on the real line, and in which two intervals are adjacent if and only if they overlap.

**Conjecture** (Stanley–Stembridge, 1993). For every indifference graph  $G$ ,  $X_G$  is e-positive.

# Indifference graphs

Note that in the above example,  $X_G$  is **e-positive**; i.e., it is a polynomial in the  $e_i$  with **nonnegative** coefficients.

This is not true for all graphs; however. . .

**Definition.** An **indifference graph** is a graph whose vertex set is a set of (distinct) closed unit intervals on the real line, and in which two intervals are adjacent if and only if they overlap.

**Conjecture** (Stanley–Stembridge, 1993). For every indifference graph  $G$ ,  $X_G$  is e-positive.

**Note.** The original Stanley–Stembridge conjecture was seemingly more general; Guay-Paquet reduced it to the statement above.

## Schur function expansion of $X_G$

**Theorem** (Haiman 1993, Gasharov 1996). If  $G$  is an indifference graph then  $X_G$  is **s-positive**, i.e., a *nonnegative* linear combination of Schur functions. The coefficients count certain tableau-like objects.

## Schur function expansion of $X_G$

**Theorem** (Haiman 1993, Gasharov 1996). If  $G$  is an indifference graph then  $X_G$  is **s-positive**, i.e., a *nonnegative* linear combination of Schur functions. The coefficients count certain tableau-like objects.

**Question.** If  $G$  is an indifference graph, is  $X_G = \text{ch } \rho$  for some “naturally occurring” representation  $\rho$ , where  $\text{ch}$  denotes the characteristic map?

## Schur function expansion of $X_G$

**Theorem** (Haiman 1993, Gasharov 1996). If  $G$  is an indifference graph then  $X_G$  is **s-positive**, i.e., a *nonnegative* linear combination of Schur functions. The coefficients count certain tableau-like objects.

**Question.** If  $G$  is an indifference graph, is  $X_G = \text{ch } \rho$  for some “naturally occurring” representation  $\rho$ , where  $\text{ch}$  denotes the characteristic map?

**Conjecture** (Shareshian–Wachs, 2011).  $\text{ch } \rho = \omega X_G$  where  $\rho$  is the dot action on the cohomology of a regular semisimple Hessenberg variety and  $\omega$  is a standard involution on symmetric functions. In fact,  $\omega X_G$  naturally decomposes into summands corresponding to each cohomology group  $H^{2d}$  separately.

## Schur function expansion of $X_G$

**Theorem** (Haiman 1993, Gasharov 1996). If  $G$  is an indifference graph then  $X_G$  is **s-positive**, i.e., a *nonnegative* linear combination of Schur functions. The coefficients count certain tableau-like objects.

**Question.** If  $G$  is an indifference graph, is  $X_G = \text{ch } \rho$  for some “naturally occurring” representation  $\rho$ , where  $\text{ch}$  denotes the characteristic map?

**Conjecture** (Shareshian–Wachs, 2011).  $\text{ch } \rho = \omega X_G$  where  $\rho$  is the dot action on the cohomology of a regular semisimple Hessenberg variety and  $\omega$  is a standard involution on symmetric functions. In fact,  $\omega X_G$  naturally decomposes into summands corresponding to each cohomology group  $H^{2d}$  separately.

Our main result is a proof of this conjecture (2015). Shortly afterwards, Guay-Paquet gave an independent proof using completely different methods (Hopf algebras).

# Classification of indifference graphs

Let  $\mathbf{m} = (m_1, \dots, m_{n-1})$  be a weakly increasing sequence of integers such that  $i \leq m_i \leq n$  for all  $i$ .

**Example.** If  $n = 3$  then  $\mathbf{m} \in \{(1, 2), (1, 3), (2, 2), (2, 3), (3, 3)\}$ .

# Classification of indifference graphs

Let  $\mathbf{m} = (m_1, \dots, m_{n-1})$  be a weakly increasing sequence of integers such that  $i \leq m_i \leq n$  for all  $i$ .

**Example.** If  $n = 3$  then  $\mathbf{m} \in \{(1, 2), (1, 3), (2, 2), (2, 3), (3, 3)\}$ .

Let  $G(\mathbf{m})$  be the graph with vertex set  $\{1, 2, \dots, n\}$  and in which  $i$  and  $j$  are adjacent if  $i < j \leq m_i$ .

**Example.** If  $m_i = i + 1$  for all  $i$  then  $G(\mathbf{m})$  is a path.

# Classification of indifference graphs

Let  $\mathbf{m} = (m_1, \dots, m_{n-1})$  be a weakly increasing sequence of integers such that  $i \leq m_i \leq n$  for all  $i$ .

**Example.** If  $n = 3$  then  $\mathbf{m} \in \{(1, 2), (1, 3), (2, 2), (2, 3), (3, 3)\}$ .

Let  $G(\mathbf{m})$  be the graph with vertex set  $\{1, 2, \dots, n\}$  and in which  $i$  and  $j$  are adjacent if  $i < j \leq m_i$ .

**Example.** If  $m_i = i + 1$  for all  $i$  then  $G(\mathbf{m})$  is a path.

**Fact** (implicit in the literature, explicit in Shareshian–Wachs).  
 $G(\mathbf{m})$  is an indifference graph, and every indifference graph is isomorphic to some  $G(\mathbf{m})$ .

## Hessenberg varieties

A **complete flag** in an  $n$ -dimensional vector space  $V$  is a nested sequence of subspaces  $F_1 \subseteq F_2 \subseteq \cdots \subseteq F_n = V$  such that  $\dim F_i = i$  for all  $i$ . The set of all complete flags forms a space called the **complete flag variety**.

## Hessenberg varieties

A **complete flag** in an  $n$ -dimensional vector space  $V$  is a nested sequence of subspaces  $F_1 \subseteq F_2 \subseteq \cdots \subseteq F_n = V$  such that  $\dim F_i = i$  for all  $i$ . The set of all complete flags forms a space called the **complete flag variety**.

Let  $\mathbf{m} = (m_1, \dots, m_{n-1})$  be a weakly increasing sequence of integers such that  $i \leq m_i \leq n$  for all  $i$ .

## Hessenberg varieties

A **complete flag** in an  $n$ -dimensional vector space  $V$  is a nested sequence of subspaces  $F_1 \subseteq F_2 \subseteq \cdots \subseteq F_n = V$  such that  $\dim F_i = i$  for all  $i$ . The set of all complete flags forms a space called the **complete flag variety**.

Let  $\mathbf{m} = (m_1, \dots, m_{n-1})$  be a weakly increasing sequence of integers such that  $i \leq m_i \leq n$  for all  $i$ .

**Definition** (De Mari–Procesi–Shayman). Let  $s$  be an  $n \times n$  matrix. The **Hessenberg variety**  $\mathcal{H}(\mathbf{m}, s)$  is defined by

$$\mathcal{H}(\mathbf{m}, s) := \{\text{complete flags such that } sF_i \subseteq F_{m_i} \text{ for all } i.\}$$

If  $s$  is diagonalizable, we say  $\mathcal{H}(\mathbf{m}, s)$  is **semisimple**. If the Jordan blocks of  $s$  have distinct eigenvalues, we say  $\mathcal{H}(\mathbf{m}, s)$  is **regular**.

# The dot action

Diagonal matrices form a torus  $T$  that acts on  $\mathcal{H}(\mathbf{m}, s)$ .

Hessenberg varieties have no odd-dimensional cohomology, so in particular, **Goresky–Kottwitz–MacPherson** theory tells us that the  $T$ -equivariant cohomology can be completely described by a combinatorial object called the **moment graph**.

The vertices of the moment graph are the  $T$ -fixed points and the edges are the one-dimensional  $T$ -orbits.

# The dot action

Diagonal matrices form a torus  $T$  that acts on  $\mathcal{H}(\mathbf{m}, s)$ .

Hessenberg varieties have no odd-dimensional cohomology, so in particular, **Goresky–Kottwitz–MacPherson** theory tells us that the  $T$ -equivariant cohomology can be completely described by a combinatorial object called the **moment graph**.

The vertices of the moment graph are the  $T$ -fixed points and the edges are the one-dimensional  $T$ -orbits.

Ordinary cohomology is a quotient of equivariant cohomology. Tymoczko defined an action, the **dot action**, of the symmetric group on the cohomology of a regular semisimple Hessenberg variety  $\mathcal{H}(\mathbf{m}, s)$ . The action depends only on  $\mathbf{m}$  and not on the choice of regular semisimple  $s$ .

## Linchpin of proof

**Theorem.** Let  $\lambda$  be a partition of  $n$ . Let  $S_\lambda := S_{\lambda_1} \times \cdots \times S_{\lambda_\ell}$  be a Young subgroup of  $S_n$ . Let  $s$  be a regular matrix with Jordan type  $\lambda$ . Then the dimension of the subspace of  $H^{2d}$  fixed by  $S_\lambda$  under the dot action on a regular *semisimple* Hessenberg variety equals the Betti number  $\beta_{2d}$  of  $\mathcal{H}(\mathbf{m}, s)$ .

## Linchpin of proof

**Theorem.** Let  $\lambda$  be a partition of  $n$ . Let  $S_\lambda := S_{\lambda_1} \times \cdots \times S_{\lambda_\ell}$  be a Young subgroup of  $S_n$ . Let  $s$  be a regular matrix with Jordan type  $\lambda$ . Then the dimension of the subspace of  $H^{2d}$  fixed by  $S_\lambda$  under the dot action on a regular *semisimple* Hessenberg variety equals the Betti number  $\beta_{2d}$  of  $\mathcal{H}(\mathbf{m}, s)$ .

**Standard fact.** The dimensions of the above fixed subspaces are the coefficients of the **monomial symmetric function** expansion.

Therefore the above theorem reduces the computation of the dot action to the computation of regular (but not necessarily semisimple) Hessenberg varieties.

## The combinatorial part of the proof

Tymoczko has already obtained a combinatorial description of the cohomology of  $\mathcal{H}(\mathbf{m}, s)$  for all  $s$ .

# The combinatorial part of the proof

Tymoczko has already obtained a combinatorial description of the cohomology of  $\mathcal{H}(\mathbf{m}, s)$  for all  $s$ .

Connecting Tymoczko's work to  $X_G$  takes two steps.

1. We generalize a **combinatorial reciprocity theorem** of Chow to obtain a combinatorial description of the coefficients in the monomial symmetric function expansion.

# The combinatorial part of the proof

Tymoczko has already obtained a combinatorial description of the cohomology of  $\mathcal{H}(\mathbf{m}, s)$  for all  $s$ .

Connecting Tymoczko's work to  $X_G$  takes two steps.

1. We generalize a **combinatorial reciprocity theorem** of Chow to obtain a combinatorial description of the coefficients in the monomial symmetric function expansion.
2. We describe an explicit bijection between our combinatorial description and Tymoczko's combinatorial description when  $s$  is regular.

# The combinatorial part of the proof

Tymoczko has already obtained a combinatorial description of the cohomology of  $\mathcal{H}(\mathbf{m}, s)$  for all  $s$ .

Connecting Tymoczko's work to  $X_G$  takes two steps.

1. We generalize a **combinatorial reciprocity theorem** of Chow to obtain a combinatorial description of the coefficients in the monomial symmetric function expansion.
2. We describe an explicit bijection between our combinatorial description and Tymoczko's combinatorial description when  $s$  is regular.

**Corollary.** The Betti numbers of a regular Hessenberg variety form a palindromic sequence. (Follows from a theorem of Shareshian and Wachs. Note that regular Hessenberg varieties are not smooth, and the corollary is not true if  $s$  is not regular. This corollary has since been generalized to other types by Precup.)

# The geometric part of the proof

A monodromy argument relates the  $S_\lambda$  invariants to a space of **local invariant cycles**.

Work of Beilinson–Bernstein–Deligne on perverse sheaves then implies that there is a **surjection** from the cohomology of regular Hessenberg varieties to the space of local invariant cycles.

# The geometric part of the proof

A monodromy argument relates the  $S_\lambda$  invariants to a space of **local invariant cycles**.

Work of Beilinson–Bernstein–Deligne on perverse sheaves then implies that there is a **surjection** from the cohomology of regular Hessenberg varieties to the space of local invariant cycles.

We then show that the palindromicity of the Betti numbers implies that the surjection is actually an isomorphism.

# The geometric part of the proof

A monodromy argument relates the  $S_\lambda$  invariants to a space of **local invariant cycles**.

Work of Beilinson–Bernstein–Deligne on perverse sheaves then implies that there is a **surjection** from the cohomology of regular Hessenberg varieties to the space of local invariant cycles.

We then show that the palindromicity of the Betti numbers implies that the surjection is actually an isomorphism.


**Note.** Abe–Harada–Horiguchi–Masuda previously carried out a similar argument in the special case of regular nilpotent  $s$ .


## BACKUP SLIDES

## The moment graph

Example on right:  $n = 3$ ,  $m_i = i + 1$ .

- ▶ The vertices are the permutations of  $\{1, 2, \dots, n\}$ .
- ▶ A transposition  $(i, j)$  is **admissible** if  $i < j \leq m_i$ . For  $m_i = i + 1$ , these are the adjacent transpositions.
- ▶ Two permutations are adjacent if they differ by an admissible transposition on **positions**.
- ▶ An edge is labeled with  $t_i - t_j$  where  $i$  and  $j$  are the transposed **numbers**.





## The dot action

- ▶ An **equivariant cohomology class**  $c$  is an assignment of a polynomial  $c(w)$  in the  $t$ 's to each vertex  $w$  such that polynomials on adjacent vertices differ by a multiple of the edge label.

# The dot action

- ▶ An **equivariant cohomology class**  $c$  is an assignment of a polynomial  $c(w)$  in the  $t$ 's to each vertex  $w$  such that polynomials on adjacent vertices differ by a multiple of the edge label.
- ▶ If  $\sigma \in S_n$  then  $(\sigma c)(w)$  is obtained by taking  $c(\sigma^{-1}w)$  (where  $\sigma^{-1}w$  means letting  $\sigma^{-1}$  act on the *numbers* of  $w$ ) and then applying  $\sigma$  to the *subscripts* of the  $t$ 's.

# The dot action

- ▶ An **equivariant cohomology class**  $c$  is an assignment of a polynomial  $c(w)$  in the  $t$ 's to each vertex  $w$  such that polynomials on adjacent vertices differ by a multiple of the edge label.
- ▶ If  $\sigma \in S_n$  then  $(\sigma c)(w)$  is obtained by taking  $c(\sigma^{-1}w)$  (where  $\sigma^{-1}w$  means letting  $\sigma^{-1}$  act on the *numbers* of  $w$ ) and then applying  $\sigma$  to the *subscripts* of the  $t$ 's.
- ▶ Equivariant cohomology classes comprise a free module over  $\mathbb{C}[t_1, \dots, t_n]$ . Write down matrices for the above representation with respect to some basis, and then take the constant terms of the entries to get the dot action on the cohomology.